

# Assessment of Blood Pressure Monitoring During the Postpartum Period

Kamila Drobina<sup>1</sup>, Anna Prokopowicz<sup>1</sup>, Izabella Uchmanowicz<sup>2</sup>

<sup>1</sup> Division of Fundamentals of Midwifery, Department of Midwifery, Faculty of Nursing and Midwifery, Wrocław Medical University, Wrocław, Poland

<sup>2</sup> Division of Research Methodology, Department of Nursing, Faculty of Nursing and Midwifery, Wrocław Medical University, Wrocław, Poland

(Corresponding Author: Kamila Drobina, [kamila.drobina@umw.edu.pl](mailto:kamila.drobina@umw.edu.pl))

**Citation:** Drobina, K., Prokopowicz, A., & Uchmanowicz, I. (2026). Assessment of blood pressure monitoring during the postpartum period. *International Journal of Health Supplements*, 1(Suppl. 3), 316–317. <https://doi.org/10.82051/ijhs-2026-Suppl.3>

**Introduction.** Pregnancy-induced hypertension affects 5–10% of women, while approximately 9% may develop de novo postpartum hypertension. Blood pressure monitoring is essential for early detection of abnormalities and cardiovascular complications. This study aimed to assess postpartum self-monitoring, blood pressure measurements performed by healthcare professionals, and recommendations provided at hospital discharge.

**Methods.** This pilot analysis was part of an ongoing longitudinal study investigating pregnancy-induced hypertension and women's emotional state. Among 104 postpartum women recruited at a tertiary referral centre in Wrocław, 70 completed follow-up assessments. Data were collected on postpartum days 2–3 and at 5–6 weeks postpartum using standardised questionnaires and an author-developed survey.

**Results.** All participants monitored blood pressure during pregnancy, compared with 50% (n=35) postpartum. In hospital, 87% (n=61) received daily measurements, while 10% (n=7) received none. Outpatient monitoring was reported during gynaecological visits by 49% (n=32) and midwife visits by 17% (n=12). Among women with pregnancy-related hypertension (n=12), 75% (n=9) performed postpartum self-monitoring. Their motivation was higher than that of normotensive women (median 9 vs 1). Overall, 20% (n=14) received discharge recommendations for home monitoring, including 67% (n=8) with pregnancy-related hypertension and 11% (n=6) without hypertension.

**Discussion.** Women with pregnancy-related hypertension demonstrated greater motivation for postpartum blood pressure monitoring and received more education from healthcare professionals. Education among women without hypertension was insufficient, potentially limiting preventive self-monitoring. Larger studies are needed to evaluate whether systematic discharge education improves postpartum blood pressure monitoring.

**Keywords:** Postpartum Period, Blood Pressure Monitoring, Hypertension, Health Education, Hypertensive Disorders of Pregnancy

---

**Funding:** This research received no external funding. **Conflict of Interest:** The authors declare no conflict of interest.