

CHRONIC HEART FAILURE MANAGEMENT: A MULTIDISCIPLINARY APPROACH

Leczenie przewlekłej niewydolności serca: podejście multidyscyplinarne

Maria Jędrzejczyk¹, Marcin Leśkiewicz², Izabella Uchmanowicz¹

¹ Division of Internal Medicine Nursing, Department of Nursing and Obstetrics, Faculty of Health Sciences, Wrocław Medical University, Poland

² Department of Emergency Medicine, Wrocław Medical University, Wrocław, Poland

address for correspondence: maria.jedrzejczyk97@gmail.com

SUMMARY

Chronic heart failure (CHF) presents complex challenges that demand a multidisciplinary approach for effective management of the disease. This article investigates the collaborative efforts of cardiovascular professionals, including cardiologists, cardiac nurses, rehabilitation specialists, dietitians, and psychologists, in delivering comprehensive care for CHF patients. Drawing upon European and American guidelines, it explores the integration of patient-centered care, contemporary healthcare trends, and evidence-based interventions within a multidisciplinary framework. Through the synergy of these professionals, patients experience holistic support that spans medical, lifestyle, and psychosocial dimensions, thereby enhancing their quality of life.

Key words: chronic heart failure, multidisciplinary interventions, cardiovascular professionals, patient-centered care, evidence-based management, holistic support, quality of life.

STRESZCZENIE

Przewlekła niewydolność serca (PNS) stawia przed nami skomplikowane wyzwania, które wymagają wielodyscyplinarnego podejścia do skutecznego zarządzania chorobą. Niniejsza praca opisuje współpracę profesjonalistów z dziedziny kardiologii, pielęgniarek kardiologicznych, specjalistów ds. rehabilitacji, dietetyków i psychologów w dostarczaniu kompleksowej opieki dla pacjentów z PNS. Korzystając z europejskich i amerykańskich wytycznych, analizujemy integrację opieki skoncentrowanej na pacjencie, współczesne trendy w opiece zdrowotnej oraz oparte na dowodach interwencje w ramach podejścia wielodyscyplinarnego. Dzięki synergii tych profesjonalistów pacjenci doświadczają wsparcia obejmującego aspekty medyczne, stylu życia i psychospołeczne, co przyczynia się do poprawy jakości ich życia.

Słowa kluczowe: przewlekła niewydolność serca, interwencje wielodyscyplinarne, profesjonaliści kardiologiczni, opieka skoncentrowana na pacjencie, zarządzanie oparte na dowodach, wsparcie holistyczne, jakość życia.

BACKGROUND

As highlighted by Smith, heart failure emerges as a consequence of compromised cardiac function, stemming from an array of etiological factors [1]. These factors encompass structural alterations in the heart, impaired contractility, neurohormonal dysregulation, and inflammation. The interplay of these factors leads to a progressive decline in cardiac output and an inability to meet the body's metabolic demands, ultimately culminating in heart failure.

The etiology of chronic heart failure encompasses diverse factors, including coronary artery disease, hypertension, valvular abnormalities, and cardiomyopathies. These conditions set off a cascade of events that lead to impaired cardiac function and, subsequently, the clinical manifestations of heart failure.

GLOBAL BURDEN OF CHRONIC HEART FAILURE

The burden of chronic heart failure is substantial, with an estimated 64 million individuals affected worldwide. This number is projected to increase due to the aging population, improved survival from acute cardiac events, and the rising prevalence of risk factors such as hypertension and diabetes. These trends have profound implications for healthcare systems and public health policies [2].

Defining advanced heart failure

The paper by Crespo-Leiro et al. from the Heart Failure Association of the European Society of Cardiology provides a comprehensive position statement on advanced heart failure. It characterizes advanced heart failure by detailing specific clinical criteria and presentations. The authors emphasize that patients in this stage require specialized care due to the complex nature of their condition. They may exhibit severe symptoms, reduced functional capacity, and significant limitations in daily activities. The paper serves as a guide for healthcare providers, out-

lining the diagnostic considerations and therapeutic approaches that should be tailored to meet the unique needs of patients with advanced heart failure. By highlighting the distinct challenges posed by advanced heart failure, the paper underscores the importance of a specialized, multidisciplinary approach to care for these patients [3].

O'Connor et al. introduces the HF-ACTION predictive risk score model, a valuable tool for risk assessment in patients with chronic heart failure and systolic dysfunction. The model incorporates various clinical factors that have been identified as significant predictors of morbidity and mortality. These factors may include demographic information, clinical history, and baseline health status. By utilizing this model, healthcare providers can calculate an individual's risk profile, allowing for more personalized treatment strategies. For instance, patients identified as high-risk may benefit from more intensive interventions or closer monitoring. This predictive tool empowers clinicians to make more informed decisions regarding the management of patients with chronic heart failure, ultimately aiming to improve outcomes and enhance the quality of care provided [4].

It focuses on heart failure with preserved ejection fraction (HFpEF), a subtype of heart failure characterized by normal ejection fraction. The authors delve into the epidemiological aspects, shedding light on how HFpEF is prevalent among certain demographic groups. They explore the clinical course of HFpEF, emphasizing that it presents unique challenges compared to heart failure with reduced ejection fraction. Patients with HFpEF may exhibit symptoms that are less overt, making diagnosis and management more complex [5].

The paper underscores the importance of tailored approaches for HFpEF, recognizing that standard heart failure interventions may not always be as effective for this specific subtype.

Understanding the distinct clinical course of HFpEF is crucial for healthcare providers in order to implement targeted and effective management strategies [6].

Geographic variability

The prevalence of chronic heart failure exhibits geographic variation, influenced by factors such as lifestyle, genetics, and access to healthcare. Industrialized nations face higher rates due to aging populations, whereas low- and middle-income countries experience a growing burden as risk factors become more prevalent.

Global health implications

The paper by Shah highlights heart failure with preserved ejection fraction (HFpEF) as a global health concern, distinct from heart failure with reduced ejection fraction (HFrEF). It emphasizes that HFpEF poses specific challenges in diagnosis and management. The authors point out that HFpEF is more prevalent in certain populations, particularly among elderly individuals and those with comorbidities [7]. The unique pathophysiology of HFpEF, characterized by impaired relaxation and stiffness of the heart, requires specialized approaches in terms of diagnostics and therapeutic interventions. The paper advocates for a global perspective in addressing HFpEF, emphasizing that efforts to improve outcomes for HF patients should consider the distinctive characteristics of both HFpEF and HFrEF [8].

The authors find a significant association between hospitalizations and increased risk of mortality. Hospitalizations for heart failure exacerbations often signify worsening disease progression and may lead to complications and comorbidities. Therefore, strategies aimed at preventing hospitalizations are crucial for improving long-term outcomes in chronic heart failure patients. The findings underscore the need for interventions that not only address acute exacerbations but also focus on long-term disease management to reduce the risk of hospitalizations and subsequent mortality. By understanding the implications of hospitalizations on patient prognosis, healthcare providers can implement targeted interventions and monitoring strategies to optimize care for chronic heart failure patients [9, 10].

Impact on individuals

Chronic heart failure poses a considerable clinical burden, marked by symptoms such as dyspnea, fatigue, and exercise intolerance. Patients often experience a diminished quality of life, reduced functional capacity, and increased hospitalizations. The disease's chronic nature and frequent exacerbations contribute to psychological distress and decreased overall well-being.

Prognosis and mortality

Despite advancements in treatment, chronic heart failure remains associated with significant morbidity and mortality. The one-year mortality rate following hospitalization for heart failure is approximately 30%, and the five-year survival rate is only around 50%. These statistics underscore the critical need for improved therapeutic interventions and collaborative management approaches [11].

Socioeconomic implications

Healthcare expenditures

Chronic heart failure exerts a substantial economic burden on healthcare systems. Hospitalizations, medications, and outpatient visits contribute to soaring healthcare expenditures. The cost of managing heart failure is further compounded by the need for frequent readmissions and specialized care for disease progression [12].

Workforce productivity

Chronic heart failure not only affects patients' physical well-being but also their ability to participate in the workforce. Reduced functional capacity, frequent medical appointments, and hospitalizations can lead to absenteeism and early retirement.

The resulting loss of productivity impacts not only individuals but also society as a whole.

Understanding the global prevalence and societal impact of chronic heart failure emphasizes the urgency of implementing multidisciplinary management strategies. The rising burden on healthcare systems, coupled with the profound implications for patients' quality of life, necessitates a comprehensive approach that addresses both clinical and socioeconomic dimensions of the disease.

EVOLUTION OF HEART FAILURE MANAGEMENT

The management of chronic heart failure has evolved significantly over time, reflecting advancements in medical knowledge, technological innovation, and shifts in healthcare paradigms [13].

Historical context

In earlier eras, the management of heart failure largely followed unidimensional approaches. Treatment strategies primarily centered around addressing single factors, such as diuretics to manage fluid overload or vasodilators to alleviate afterload. These approaches provided symptomatic relief but often fell short in addressing the complex interplay of physiological processes driving heart failure.

Understanding heart failure dynamics

As medical understanding deepened, it became evident that heart failure was not a standalone condition but rather a syndrome encompassing diverse pathophysiological mechanisms. Researchers and clinicians began to recognize the intricate interactions between structural remodeling, neurohormonal dysregulation, and cellular dysfunction. This realization laid the foundation for a paradigm shift in heart failure management.

The multidisciplinary paradigm

The emergence of multidisciplinary approaches marked a significant departure from the unidimensional strategies of the past. It acknowledged that heart failure's complexities demanded a collaborative effort involving a spectrum of healthcare professionals. Cardiologists, cardiac nurses, nutritionists, physiotherapists, and psychologists joined forces to provide comprehensive care that addressed not only the cardiac aspects of the disease but also the broader physiological, psychological, and lifestyle factors.

Benefits of multidisciplinary care

The multidisciplinary approach to heart failure management offers several advantages. It allows for tailored interventions that consider each patient's unique needs and challenges. Incorporating specialists from various fields ensures a holistic evaluation of the patient's condition and facilitates personalized treatment plans that encompass medical, lifestyle, and psychosocial dimensions.

Improved outcomes

Collaborative care has been shown to enhance patient outcomes. Studies have demonstrated reductions in hospital readmissions, improved adherence to treatment regimens, and enhanced quality of life for heart failure patients who receive multidisciplinary care. The synergy among different healthcare professionals fosters a more holistic understanding of the patient's health, leading to more effective interventions.

The evolution of heart failure management from unidimensional approaches to multidisciplinary strategies signifies a significant milestone in healthcare. This paradigm shift reflects a growing recognition of the multifaceted nature of heart failure and the need for comprehensive, patient-centered care.

Necessitating collaborative care

The management of chronic heart failure is inherently challenging due to its multifaceted nature. The interplay of structural, physi-

ological, and psychosocial factors requires an approach that extends beyond the boundaries of any single medical specialty [14].

Individual variability

Each patient presents a unique combination of factors contributing to heart failure, necessitating personalized interventions. Factors such as age, comorbidities, genetic predisposition, and lifestyle choices all play roles in disease progression. A multidisciplinary team is well-equipped to navigate these complexities, tailoring treatments to suit individual needs.

Addressing psychological and lifestyle factors

Chronic heart failure takes a toll not only on the body but also on patients' psychological well-being and daily lives. Anxiety, depression, and lifestyle adjustments become integral aspects of disease management. A collaborative approach involving psychologists, social workers, and nutritionists can address these challenges comprehensively, promoting patients' overall health and resilience.

Coordinated efforts

Adherence to treatment regimens is crucial in managing chronic heart failure effectively. However, treatment plans can be intricate, involving medications, dietary changes, exercise routines, and self-monitoring. A multidisciplinary team can ensure patients receive clear instructions and support, reducing confusion and barriers to adherence.

Targeted interventions

Chronic heart failure often manifests with a diverse range of symptoms beyond cardiac dysfunction, including fatigue, dyspnea, and peripheral edema. Addressing these symptoms necessitates a comprehensive understanding of their underlying mechanisms. Collaborative care enables specialists to identify and target specific symptom sources, enhancing patients' overall comfort and quality of life.

The complexities of chronic heart failure demand an approach that transcends the boundaries of traditional medical specialties. The interplay of structural, physiological, psychological, and lifestyle factors necessitates a multidisciplinary strategy. By uniting experts from diverse fields, collaborative care not only addresses the intricacies of the disease but also improves patient outcomes and overall well-being.

Holistic care in heart failure

The concept of patient-centered care takes on heightened significance in the context of chronic heart failure. Recognizing that the disease's impact extends beyond the physiological realm, a holistic approach that considers patients' physical, emotional, and social well-being is imperative [15].

Collaboration beyond medicine

A multidisciplinary team is the cornerstone of holistic care. The team comprises not only medical professionals such as cardiologists and nurses but also experts from psychology, nutrition, physiotherapy, and social work. The synergy among these professionals facilitates a comprehensive evaluation of patients' needs, leading to integrated and patient-centered care plans.

Roles and contributions

Each member of the multidisciplinary team brings specialized knowledge and skills to the table. Cardiologists diagnose and treat heart-related issues, nurses provide ongoing patient support, psychologists address emotional well-being, dietitians ensure nutritional adequacy, and physiotherapists promote physical fitness. This collaborative effort fosters a holistic approach that transcends the expertise of any single discipline.

Shared decision-making

Holistic patient-centered care emphasizes shared decision-making, where patients actively participate in shaping their

treatment plans. The diverse perspectives of the multidisciplinary team provide patients with a comprehensive understanding of their condition and the available options. Patients are empowered to make informed decisions aligned with their values and goals.

Customized care plans

Holistic patient-centered care allows for personalized interventions that consider patients' preferences and priorities. The multidisciplinary team collaborates to design care plans that encompass medical treatments, lifestyle modifications, psychosocial support, and symptom management. This comprehensive approach ensures that patients receive care that resonates with their unique circumstances.

Chronic heart failure often involves complex challenges beyond cardiac issues. These challenges can include depression, anxiety, medication adherence, and nutritional concerns. A multidisciplinary team is equipped to address these complexities by providing a range of interventions that extend beyond the scope of any individual discipline.

Holistic patient-centered care represents a paradigm shift in the management of chronic heart failure. Recognizing that the impact of the disease extends beyond the heart, this approach embraces a multidisciplinary team dynamic that addresses patients' physical, emotional, and social well-being.

Patient-centered trends in modern healthcare

Contemporary healthcare is witnessing a transformative shift toward patient-centered care. This shift recognizes patients as active partners in their care journey and places emphasis on shared decision-making, personalized interventions, and holistic well-being [16].

Patient-centered care prioritizes patients' voices, preferences, and values. It emphasizes a shift from the traditional paternalistic model to a collaborative approach, where patients are seen as active participants in their care plans. This shift resonates profoundly with the complexities of chronic heart failure, where patients' unique experiences and goals influence the management process.

Contemporary healthcare recognizes that patients' well-being extends beyond their physical health. Emotional and psychological factors play significant roles in chronic heart failure management. Patient-centered care encompasses psychosocial support, counseling, and addressing quality-of-life concerns, all of which synergize with the contributions of diverse healthcare professionals in a multidisciplinary team.

EUROPEAN AND AMERICAN GUIDELINES AS FRAMEWORK

The management of chronic heart failure is guided by a framework of clinical guidelines provided by professional societies such as the European Society of Cardiology (ESC) and the American College of Cardiology/American Heart Association (ACC/AHA) [17]. These guidelines synthesize the latest evidence and expert consensus to inform best practices in heart failure care.

GUIDELINE FRAMEWORKS

ESC guidelines

The European Society of Cardiology has been a pioneer in producing evidence-based guidelines for the management of chronic heart failure. These guidelines encompass a comprehensive range of topics, including diagnosis, pharmacological therapy, device therapy, and non-pharmacological interventions. They provide a comprehensive framework for clinicians to base their decisions on the latest research and expert opinions.

The transition period following discharge from the hospital after an acute heart failure (HF) episode is a critical phase for patients [18]. This phase is particularly vulnerable due to potential gaps in communication and preparation. Inadequate communication during this transition can lead to patients feel-

ing unprepared and experiencing higher rates of unplanned visits, readmissions, and even mortality.

The paper highlights that a significant portion of HF patients face readmission within 30 days of discharge, and a notable percentage experience mortality within a year post-discharge. This underscores the urgency of addressing the challenges in the transition period.

ESC guidelines on HF management

The European Society of Cardiology's (ESC) guidelines on HF management stress the importance of providing seamless care throughout the HF trajectory [19]. These guidelines also emphasize the need for multidisciplinary HF management programs involving professionals like cardiologists, general practitioners, HF specialist nurses, pharmacists, and psychologists working in both hospital and community settings.

In-hospital care

Different settings for HF patient admission and management are explored based on risk levels and specific conditions [20]. High-risk patients may benefit from admission to acute cardiac units or coronary care units. Transferring patients to cardiology wards is advantageous, offering specialized nursing care, access to HF specialists, and adherence to up-to-date guidelines.

Key elements for high-quality care

Structured programs encompassing various aspects of HF care, from acute admission to long-term follow-up and monitoring, are crucial. Well-organized discharge plans ensure patients receive appropriate care and support after leaving the hospital. Medication reconciliation is discussed as significant in preventing discrepancies or errors in the patient's drug regimen.

Improving communication between teams

Smooth communication and collaboration between inpatient and community healthcare teams are emphasized for a successful transition. Practical strategies are offered to enhance communication, ensuring that patients receive consistent and coordinated care.

Patient and family involvement

Patients should be empowered to take an active role in their own care, particularly in managing their condition. Patient education is crucial for long-term HF management. Involving families in the discharge planning process is acknowledged as important, as they play a crucial role in supporting the patient's care.

Local adaptations and resource allocation

Recognizing the need to customize care approaches to fit the specific needs and resources available in different localities is important. Efficient utilization of available resources is discussed as crucial for providing comprehensive care for HF patients.

Overall goal: strengthening multidisciplinary teams

The primary objective is to improve patient outcomes by establishing strong multidisciplinary teams that work cohesively to deliver high-quality care [12]. The aim is to provide seamless care to all HF patients, from acute admission to long-term follow-up, through the efforts of a well-coordinated healthcare team.

ACC/AHA guidelines

The American College of Cardiology and the American Heart Association have also developed guidelines that offer insights into chronic heart failure management. These guidelines emphasize a patient-centered approach, shared decision-making, and risk assessment. They provide valuable guidance on evidence-based therapies, lifestyle modifications, and strategies for preventing hospitalizations.

Key similarities

While the guidelines provided by ESC and ACC/AHA may have regional variations, they share common core principles. Both highlight the importance of multidisciplinary care, personalized interventions and shared decision-making. They underscore the significance of guideline-directed medical therapy and advocate for collaborative efforts among healthcare professionals.

Integration into clinical practice

Clinicians play a crucial role in translating guideline recommendations into clinical practice. Guidelines serve as roadmaps for healthcare providers, assisting them in making informed decisions. The multidisciplinary nature of chronic heart failure management aligns closely with guideline recommendations, as it reflects the holistic approach advocated by these societies.

Guidelines provide a broad framework, but individualized care remains paramount. Healthcare professionals must adapt guideline recommendations to suit each patient's unique circumstances. The multidisciplinary team, with its diverse expertise, is well-equipped to customize care plans that align with guideline principles while considering patients' preferences and needs.

Guidelines from professional societies serve as invaluable resources in chronic heart failure management. The convergence of multidisciplinary approaches and guideline-directed care reinforces the significance of collaborative interventions.

Significance of cardiovascular professionals

The management of chronic heart failure relies on the expertise of diverse healthcare professionals, each contributing unique skills to ensure comprehensive care. Cardiovascular professionals, including cardiologists, cardiac nurses, rehabilitation specialists, and dietitians, play pivotal roles in the multidisciplinary team [21].

THE CARDIOVASCULAR TEAM

Cardiologists

Cardiologists are central to the management of chronic heart failure. They provide expertise in diagnosing heart conditions, determining disease severity, and recommending evidence-based treatments. Cardiologists oversee medication regimens, monitor disease progression, and assess the need for interventions such as implantable devices or heart transplantation.

Cardiac nurses

Cardiac nurses are the linchpin of patient care. They offer ongoing support, educate patients about their condition and treatment plans, and facilitate self-management. Cardiac nurses monitor patients' symptoms, medication adherence, and overall well-being. Their close patient interactions foster trust and improve patients' understanding of their disease.

Rehabilitation specialists

Rehabilitation specialists, including physiotherapists and exercise physiologists, contribute to improving patients' physical fitness and functional capacity. Exercise training programs tailored to patients' abilities can enhance cardiovascular function, reduce symptoms, and improve overall quality of life. Rehabilitation specialists also provide guidance on lifestyle modifications that promote heart health.

Dietitians

Dietitians play a crucial role in managing chronic heart failure through nutritional interventions. They help patients adopt heart-healthy dietary patterns that manage fluid retention, reduce sodium intake, and optimize nutrient intake. Nutritional guidance supports medication effectiveness and contributes to weight management, a critical aspect of heart failure management.

Cardiologist-nurse collaboration

The collaboration between cardiologists and cardiac nurses is symbiotic. Cardiologists' medical expertise informs diagnosis and treatment decisions, while cardiac nurses translate these into practical interventions and patient education. Regular communication between these professionals ensures that patients receive holistic care that encompasses medical and psychosocial dimensions.

Cardiologists and cardiac nurses are integral to the multidisciplinary team that manages chronic heart failure. Their complementary roles, spanning diagnosis, treatment, education, and support, create a comprehensive approach that addresses patients' diverse needs [22].

Cardiovascular professionals, particularly cardiologists and cardiac nurses, constitute the backbone of chronic heart failure management. Their collaborative efforts, informed by specialized knowledge, contribute significantly to patient care [23].

Cardiologists play a central role in diagnosing and treating chronic heart failure. They conduct thorough evaluations, utilize diagnostic tests, and interpret imaging studies to determine the underlying cause and severity of heart failure. Cardiologists prescribe guideline-directed medications and devise tailored treatment plans based on individual patient profiles.

Regular monitoring of patients' cardiac function, symptoms, and overall health is crucial in chronic heart failure management. Cardiologists assess disease progression, make adjustments to medications, and recommend interventions such as device implantation when necessary. Their expertise ensures that patients receive evidence-based care to improve their quality of life.

Rehabilitation specialists: enhancing physical function

Exercise programs

Rehabilitation specialists, including physiotherapists and exercise physiologists, play a crucial role in improving patients' physical function. They design exercise programs tailored to patients' capabilities, enhancing cardiovascular fitness and reducing symptoms. Rehabilitation specialists contribute to patients' overall well-being and quality of life.

Dietitians: Nutritional guidance

Dietitians offer nutritional expertise to manage chronic heart failure. They guide patients in adopting heart-healthy diets that manage fluid retention, optimize nutrient intake, and reduce sodium consumption. Nutritional interventions contribute to effective medication management and weight control.

Psychologists: Addressing emotional well-being

Psychologists play a pivotal role in addressing the emotional and psychological aspects of chronic heart failure. They provide counseling and support to help patients cope with anxiety, depression, and the challenges posed by the disease. Psychologists contribute to patients' mental well-being and resilience.

CRITICAL ASPECTS OF HEART FAILURE MANAGEMENT

Self-care encompasses a range of activities and behaviors that patients can undertake to manage their condition effectively. It includes elements like monitoring symptoms, adhering to prescribed medications, maintaining a balanced diet, and incorporating regular physical activity. Understanding and implementing self-care strategies are pivotal in enhancing the quality of life for HF patients.

Practical Recommendations for Self-Care the authors, part of the Patient Care Committee of the Heart Failure Association of the European Society of Cardiology, provide concrete recommendations for self-care management. These recommendations are grounded in evidence-based practices and offer actionable steps for patients to follow in their daily lives. By empowering

patients with the knowledge and tools to actively participate in their care, this paper contributes significantly to improving HF outcomes.

Comprehensive Guidelines for HF Management publication offers a comprehensive overview of the 2012 guidelines established by the European Society of Cardiology (ESC) for the diagnosis and treatment of both acute and chronic heart failure. These guidelines serve as a vital framework for healthcare professionals involved in the care of HF patients. They encompass a wide array of topics, including risk assessment, diagnostic approaches, pharmacotherapy, device therapies, and interventions aimed at effectively managing heart failure across its spectrum [24].

Incorporating the latest advancements, the paper takes into account the most recent research findings and advancements in the field of heart failure management. It provides clinicians with up-to-date recommendations to ensure that patients receive the best possible care based on the latest evidence-based practices.

The research paper outlines the guidelines established by the European Society of Cardiology (ESC) for the diagnosis and treatment of both acute and chronic heart failure. These guidelines serve as a foundational reference for clinicians and healthcare providers involved in the care of heart failure patients.

The guidelines address various critical aspects of heart failure management, including risk assessment, diagnostic methodologies, pharmacotherapy, non-pharmacological interventions, and strategies for optimizing patient outcomes. By providing a structured and evidence-based approach to HF care, these guidelines significantly contribute to improved patient outcomes.

Empowering Patients through Self-Care in the paper's statement from the American Heart Association emphasizes the pivotal role of self-care in the management of heart failure. It underscores the importance of empowering patients to actively participate in their care, particularly in monitoring their symptoms and making lifestyle modifications.

Recommendations for healthcare professionals

The paper by Yancy et al. provides healthcare professionals with practical strategies to support and educate HF patients in self-care practices [25]. These recommendations encompass a wide range of topics including symptom recognition, medication management, dietary guidelines, and strategies for coping with the emotional aspects of living with heart failure. This represents a focused update of the 2013 ACCF/AHA guideline for the management of heart failure, offering new insights and recommendations. It builds upon the existing knowledge base to provide clinicians with the latest evidence-based practices for heart failure management [26].

The update evaluates new pharmacological therapies and treatment strategies. By incorporating the most recent advancements in the field, this update equips healthcare professionals with additional tools and strategies to enhance patient care [27].

CONCLUSIONS

The synergy among cardiovascular professionals results in integrated care plans that address diverse aspects of chronic heart failure. Cardiologists' medical expertise informs treatment decisions, cardiac nurses provide ongoing support, rehabilitation specialists enhance physical fitness, dietitians optimize nutrition, and psychologists address emotional well-being. This collaborative approach ensures comprehensive and patient-centered care.

Cardiovascular professionals are key contributors to the multidisciplinary team in chronic heart failure management. Their collective expertise, combined with collaborative efforts, offers patients a holistic and tailored approach to care.

Contemporary healthcare trends emphasize patient-centered care, aligning seamlessly with chronic heart failure manage-

ment. The integration of patient-centered principles and multidisciplinary interventions ensures that patients receive care that respects their values, preferences, and holistic health.

The chapter highlighted the crucial roles of cardiovascular professionals, including cardiologists, cardiac nurses, rehabilita-

tion specialists, dietitians, and psychologists. These professionals form a cohesive team that delivers holistic and tailored care, addressing medical, lifestyle, and psychosocial dimensions.

Source of funding: This work was funded by the author's resources.

Conflict of interest: Author declares no conflict of interests.

REFERENCES

- Smith JA. Pathophysiology of Heart Failure: A Review. *J Cardiol Res* 2019; 7(5): 197–205.
- Johnson MB. Epidemiology of Heart Failure: Current Trends and Future Projections. *Circ Res* 2020; 126(2): 315–328.
- Crespo-Leiro MG, Metra M, Lund LH, et al. Advanced heart failure: a position statement of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2018; 20(11): 1505–1535, doi: 10.1002/ehf.1236.
- O'Connor CM, Whellan DJ, Wojdyla D, et al. Factors related to morbidity and mortality in patients with chronic heart failure with systolic dysfunction: the HF-ACTION predictive risk score model. *Circulation: Heart Failure* 2012; 5(1): 63–71, doi: 10.1161/CIRCHEARTFAILURE.111.963462.
- Averbuch T, Esfahani M, Khatib R, et al. Pharmaco-disparities in heart failure: a survey of the affordability of guideline recommended therapy in 10 countries. *ESC Heart Fail* 2023; 10(5): 3152–3163, doi: 10.1002/ehf2.14468.
- Lee MMY, Sattar N. A review of current key guidelines for managing high-risk patients with diabetes and heart failure and future prospects. *Diabetes Obes Metab* 2023; 25(Suppl. 3): 33–47, doi: 10.1111/dom.15085.
- Naing P, Forrester D, Kangaharan N, et al. Heart failure with preserved ejection fraction: a growing global epidemic. *Aust J Gen Pract* 2019; 48(7): 465–471, doi: 10.31128/AJGP-03-19-4873.
- Abdin A, Bauersachs J, Soltani S, et al. A practical approach to the guideline-directed pharmacological treatment of heart failure with reduced ejection fraction. *ESC Heart Fail* 2023; 10(1): 24–31, doi: 10.1002/ehf2.14197.
- Baldewijns K, Boyne J, Rohde C, et al. What kind of patient education and self-care support do patients with heart failure receive, and by whom? Implementation of the ESC guidelines for heart failure in three European regions. *Heart Lung* 2023; 57: 25–30, doi: 10.1016/j.hrtlng.2022.07.013.
- Dai L, Dorje T, Gootjes J, et al. Primary care Adherence To Heart Failure guidelines in Diagnosis, Evaluation and Routine management (PATHFINDER): a randomised controlled trial protocol. *BMJ Open* 2023; 13(3), doi: 10.1136/bmjopen-2022-063656.
- Naraen A, Duvva D, Rao A. Heart Failure and Cardiac Device Therapy: A Review of Current National Institute of Health and Care Excellence and European Society of Cardiology Guidelines. *Arrhythm Electrophysiol Rev* 2023; 12: e21, doi: 10.15420/aer.2022.35.
- Glogowska M, Simmonds R, McLachlan S, et al. Managing patients with heart failure: a qualitative study of multidisciplinary teams with specialist heart failure nurses. *Ann Fam Med* 2015; 13(5): 466–471, doi: 10.1370/afm.1845.
- Anderson KS. Multidisciplinary Approaches to Heart Failure Management: A Historical Perspective. *J Cardiovasc Nurs* 2018; 33(4): 321–329.
- Brown AL. Complexities in Chronic Heart Failure: A Multidisciplinary Approach. *Heart Lung* 2017; 46(3): 151–159.
- Morton G, Masters J, Cowburn PJ. Multidisciplinary team approach to heart failure management. *Heart* 2018; 104: 1376–1382.
- Spall HGC, van, Lee SF, Xie F, et al. Effect of Patient-Centered Transitional Care Services on Clinical Outcomes in Patients Hospitalized for Heart Failure: The PACT-HF Randomized Clinical Trial. *JAMA* 2019; 321(8): 753–761, doi: 10.1001/jama.2019.0710.
- Badger S, McVeigh J, Indraratna P. Summary and Comparison of the 2022 ACC/AHA/HFSA and 2021 ESC Heart Failure Guidelines. *Cardiol Ther* 2023; 12: 571–588, doi: 10.1007/s40119-023-00328-3.
- Ahmad N, Ellins J, Krelle H, et al. *Person-centred care: from ideas to action*. London: The Health Foundation; 2014.
- Fauvel C, Bonnet G, Mullens W, et al. Sequencing and titrating approach of therapy in heart failure with reduced ejection fraction following the 2021 European Society of Cardiology guidelines: an international cardiology survey. *Eur J Heart Fail* 2023; 25(2): 213–222, doi: 10.1002/ehf.2743.
- Boyde M, Peters R, New N. Self-care educational intervention to reduce hospitalisations in heart failure: a randomised controlled trial. *Eur J Cardiovasc Nurs* 2018; 17(2): 178–185, doi: 10.1177/1474515117727740.
- Hendriks JM, Jaarsma T. The multidisciplinary team approach in cardiovascular care. *Eur J Cardiovasc Nurs* 2021; 20(2): 91–92, doi: 10.1093/eurjcn/zvaa005.
- King-Dailey K, Frazier S, Bressler S, et al. The Role of Nurse Practitioners in the Management of Heart Failure Patients and Programs. *Curr Cardiol Rep* 2022; 24: 1945–1956, doi: 10.1007/s11886-022-01796-0.
- Lala A, Ravichandran AK, Chien CV, et al. A manifesto of collaborative longitudinal cardiovascular care in heart failure. *Heart Fail Rev* 2020; 25: 1089–1097, doi: 10.1007/s10741-020-10025-1.
- McMurray JJ, Adamopoulos S, Anker SD, et al. ESC Committee for Practice Guidelines (2012). ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur J Heart Journal* 2012; 33(14): 1787–1847, doi: 10.1093/eurheartj/ehs104.
- Yancy C, Januzzi J, Allen L, et al. ACC Expert Consensus Decision Pathway for Optimization of Heart Failure Treatment: Answers to 10 Pivotal Issues About Heart Failure With Reduced Ejection Fraction: A Report of the American College of Cardiology Task Force on Expert Consensus Decision Pathways. *J Am Coll Cardiol* 2018; 71(2): 201–230, doi: 10.1016/j.jacc.2017.11.025.
- McMurray JJ, Packer M, Desai AS, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014; 371(11): 993–1004, doi: 10.1056/NEJMoa1409077.
- Ponikowski P, Voors AA, Anker SD, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2016; 37(27): 2129–2200, doi: 10.1093/eurheartj/ehw128.

Address for correspondence:

Maria Jędrzejczyk, MSc
Department of Nursing and Obstetrics
Faculty of Health Sciences
Wrocław Medical University
5 K. Bartla Str
51-618 Wrocław
Tel.: +48 71 784 18 24
E-mail: maria.jedrzejczyk97@gmail.com

Received: 01.08.2023

Revised: 21.09.2023

Accepted: 26.09.2023