

SPECIAL NEEDS OF PATIENTS WITH HEART FAILURE AND FRAILTY

Specjalne potrzeby pacjentów z niewydolnością serca i kruchością

Maria Jędrzejczyk¹, Bernadetta Żółkowska²,
Karol Mirkowski², Izabella Uchmanowicz^{1,3}

¹ Division of Research Methodology, Department of Nursing, Faculty of Nursing and Midwifery, Wrocław Medical University, Wrocław, Poland

² Faculty of Medicine, Wrocław Medical University, Wrocław, Poland (student)

³ Centre for Cardiovascular Health, Edinburgh Napier University, Edinburgh, UK

adres do korespondencji: izabella.uchmanowicz@umw.edu.pl, maria.jedrzejczyk97@gmail.com

SUMMARY

Heart failure (HF) and frailty are both prevalent conditions among the elderly, often coexisting and significantly complicating patient management. This review explores the special needs of patients with both heart failure and frailty, emphasizing the unique clinical challenges posed by this dual burden. Frailty exacerbates the symptoms of heart failure, contributing to poorer outcomes, including increased mortality, frequent hospitalizations, and diminished quality of life. The review discusses the pathophysiology of both conditions, diagnostic considerations, and highlights the importance of a comprehensive, multidisciplinary approach to care. Management strategies must address both the cardiovascular and functional aspects of these patients' health, integrating pharmacologic treatments, rehabilitation, and psychosocial support. The review also emphasizes the need for early detection of frailty and the implementation of geriatric principles in heart failure management. A holistic approach that combines cardiology and geriatrics is essential for improving the prognosis and quality of life in this vulnerable population. Future research should focus on tailoring interventions for frail heart failure patients to optimize their outcomes.

Key words: heart failure, frailty, elderly patients, multidisciplinary care, quality of life.

STRESZCZENIE

Niewydolność serca (NS) oraz frailty (kruchość) to powszechne schorzenia wśród osób starszych, które często występują jednocześnie, znacznie utrudniając zarządzanie pacjentem. Niniejsza praca omawia szczególne potrzeby pacjentów z współistniejącą niewydolnością serca i frailty, podkreślając unikalne wyzwania kliniczne związane z tym podwójnym obciążeniem. Frailty pogarsza objawy niewydolności serca, prowadząc do gorszych wyników leczenia, w tym wyższej śmiertelności, częstszych hospitalizacji oraz obniżonej jakości życia. Przegląd omawia patofizjologię obu schorzeń, kwestie diagnostyczne oraz podkreśla znaczenie kompleksowego, multidyscyplinarnego podejścia do leczenia. Strategie leczenia muszą uwzględniać zarówno aspekty kardiologiczne, jak i funkcjonalne zdrowia tych pacjentów, integrując leczenie farmakologiczne, rehabilitację oraz wsparcie psychospołeczne. Praca kładzie również nacisk na potrzebę wczesnego wykrywania frailty i wdrażania zasad geriatry w leczeniu niewydolności serca. Holistyczne podejście, łączące kardiologię i geriatricę, jest kluczowe dla poprawy prognoz i jakości życia tej wrażliwej grupy pacjentów. Przyszłe badania powinny koncentrować się na dostosowaniu interwencji dla pacjentów z frailty i niewydolnością serca w celu optymalizacji ich wyników zdrowotnych.

Słowa kluczowe: niewydolność serca, osłabienie, pacjenci w podeszłym wieku, opieka multidyscyplinarna, jakość życia.

BACKGROUND

The intersection of **heart failure (HF)** and **frailty** presents a complex clinical challenge. Both conditions individually carry a significant burden in terms of morbidity, mortality, and health-care resource utilization, but when they coexist, the clinical picture becomes even more complicated. Patients with both heart failure and frailty are at an elevated risk for poor outcomes, including frequent hospitalizations, reduced functional capacity, and higher mortality rates [1]. This review explores the special needs of such patients, focusing on clinical management, diagnostic considerations, and the importance of a comprehensive, multidisciplinary approach.

UNDERSTANDING HEART FAILURE AND FRAILTY

Heart failure is a clinical syndrome characterized by the inability of the heart to pump blood efficiently to meet the body's needs, leading to symptoms such as dyspnea, fatigue, and edema. Frailty, on the other hand, refers to a state of increased vulnerability to stressors, characterized by unintentional weight loss, weakness, slow gait, low physical activity, and exhaustion [2].

When frailty and heart failure coexist, patients tend to have more severe symptoms, impaired functional status, and a significantly higher risk of adverse outcomes. Frailty is often associated with worse quality of life in HF patients, particularly as it affects their ability to cope with the physical demands of

HF management, including adherence to complex treatment regimens [3].

CLINICAL IMPLICATIONS OF COEXISTING HEART FAILURE AND FRAILTY

Increased mortality and morbidity

Several studies have shown that frailty significantly contributes to worse outcomes in heart failure. In fact, frail patients with heart failure have a higher risk of mortality compared to non-frail patients, even when adjusted for other risk factors such as age, comorbidities, and severity of heart failure [4]. Frailty contributes to this heightened risk by affecting the body's resilience to acute decompensations, hospitalizations, and interventions.

Impaired response to treatment

The frailty syndrome impairs the body's ability to respond to treatment, particularly in elderly patients. For instance, frail patients are more likely to experience adverse effects from heart failure medications, such as diuretics or ACE inhibitors, due to reduced organ function or decreased ability to metabolize drugs [5,6]. Moreover, frailty may affect the tolerance of heart failure therapies, such as beta-blockers and angiotensin-converting enzyme inhibitors, which are vital in managing the disease.

Challenges in physical rehabilitation

Frailty is often linked with muscle wasting and decreased strength, which may complicate the rehabilitation process for

heart failure patients. Exercises, mobility training, and physical rehabilitation can be more challenging, and frail patients may struggle to regain functional independence after a cardiac event [7].

DIAGNOSTIC CONSIDERATIONS

Screening for frailty in heart failure

Despite the strong association between frailty and poor outcomes in heart failure, frailty is often underdiagnosed. In routine clinical practice, patients with heart failure are frequently assessed for cardiac function (e.g., ejection fraction) but may not receive formal frailty screenings. Several tools have been developed for this purpose, including the Clinical Frailty Scale (CFS) and the Frailty Phenotype approach, which assess physical performance, strength, and endurance [8].

Comprehensive assessment

A multidisciplinary assessment that evaluates both cardiac function and frailty markers is essential. Clinicians should assess the patient's nutritional status, cognitive function, mobility, strength, and psychological health to develop a tailored management plan. In frail heart failure patients, a holistic view is necessary to address both the cardiovascular and the functional aspects of their care [5].

MANAGEMENT OF HEART FAILURE AND FRAILTY

Pharmacologic management

The pharmacologic management of heart failure in frail patients requires careful consideration of drug interactions, side effects, and dosing. For instance, frail individuals may be more sensitive to diuretics, which can lead to electrolyte imbalances or dehydration. The goal is to optimize heart failure treatment while minimizing adverse effects and maintaining functional capacity.

Medications such as angiotensin-converting enzyme inhibitors (ACE inhibitors), beta-blockers, and aldosterone antagonists are essential in the management of heart failure, but the frailty syndrome may require dose adjustments and close monitoring to prevent decompensation [9].

Non-pharmacologic management

Physical therapy is crucial for improving mobility and strength in frail patients with heart failure. Interventions such as aerobic exercise, strength training, and balance exercises can improve both cardiac outcomes and overall functional capacity. Furthermore, nutritional support plays a vital role in maintaining muscle mass and preventing weight loss, which can exacerbate frailty [10].

Psychosocial interventions, including counseling or palliative care, should be incorporated to address the mental health needs

of frail patients, as they often experience increased rates of depression and anxiety [11].

Multidisciplinary approach

A multidisciplinary care team, including cardiologists, geriatricians, physical therapists, dietitians, and mental health professionals, is essential for addressing the complex needs of these patients. Coordination between different specialties ensures a comprehensive care plan tailored to both the heart failure and frailty components [12].

PALLIATIVE AND END-OF-LIFE CARE

Given the high mortality rates and poor prognosis associated with the combination of heart failure and frailty, early palliative care is often recommended. A palliative approach focuses on symptom management, improving the quality of life, and supporting the patient's family through difficult decisions, including the potential for advanced care planning and end-of-life decisions [11].

CHALLENGES IN RESEARCH AND FUTURE DIRECTIONS

Despite growing recognition of the importance of frailty in heart failure management, there remains a lack of consensus on the best tools for frailty assessment and the optimal treatment strategies for frail heart failure patients. Most clinical trials have excluded frail individuals, and as such, evidence for treatment in this group remains limited. Future research should focus on evaluating interventions that are tailored specifically to this population, with a particular emphasis on geriatric cardiology, which aims to integrate geriatrics and cardiology for better management of elderly patients with complex comorbidities.

CONCLUSIONS

Patients with both heart failure and frailty represent a vulnerable and often neglected population. Their clinical needs are multifaceted, involving careful management of both the cardiovascular and functional aspects of their health. A thorough, multidisciplinary approach is crucial for optimizing care and improving outcomes. Further research into tailored interventions and more robust clinical trials involving frail heart failure patients is essential for improving their prognosis and quality of life.

By improving early detection of frailty and integrating geriatric principles into the management of heart failure, healthcare providers can offer a more comprehensive and personalized approach to care, ultimately improving the lives of these highly vulnerable patients.

Source of funding: This work was funded by the authors resources.

Conflict of interest: Authors declare no conflict of interests.

REFERENCES

- Vitale C, Uchmanowicz I. Frailty in patients with heart failure. *Eur Heart J Suppl* 2019; 21(Suppl. L): L12–L16, doi: 10.1093/eurheartj/suz238.
- Pandey A, Kitzman D, Reeves G. Frailty is Intertwined with Heart Failure: Mechanisms, Prevalence, Prognosis, Assessment, and Management. *JACC Heart Fail* 2019; 7(12): 1001–1011, doi: 10.1016/j.jchf.2019.10.005.
- Talha KM, Pandey A, Fudim M, et al. Frailty and heart failure: State-of-the-art review. *J Cachexia Sarcopenia Muscle* 2023; 14(5): 1959–1972, doi: 10.1002/jcsm.13306.
- Haydock PM, Flett AS. Management of heart failure with reduced ejection fraction. *Heart* 2022; 108(19): 1571–1579, doi: 10.1136/heartjnl-2020-318811.
- Afilalo J, Alexander KP, Mack MJ, et al. Frailty assessment in the cardiovascular care of older adults. *J Am Coll Cardiol* 2014; 63(8): 747–762, doi: 10.1016/j.jacc.2013.09.070.
- Wleklik M, Denfeld Q, Czaplak M, et al. A patient with heart failure, who is frail: How does this affect therapeutic decisions? *Cardiol J* 2023; 30(5): 825–831, doi: 10.5603/CJ.a2023.0027.
- Naylor M, Vasan RS. Preventing heart failure: the role of physical activity. *Curr Opin Cardiol* 2015; 30(5): 543–550, doi: 10.1097/HCO.0000000000000206.
- Lai HY, Huang ST, Anker SD, et al. The burden of frailty in heart failure: Prevalence, impacts on clinical outcomes and the role of heart failure medications. *J Cachexia Sarcopenia Muscle* 2024; 15(2): 660–670, doi: 10.1002/jcsm.13412.
- Vitale C, Spoletini I, Rosano GM. Frailty in Heart Failure: Implications for Management. *Card Fail Rev* 2018; 4(2): 104–106, doi: 10.15420/cfr.2018.22.2.
- Angulo J, El Assar M, Álvarez-Bustos A, et al. Physical activity and exercise: Strategies to manage frailty. *Redox Biol* 2020; 35: 101513, doi: 10.1016/j.redox.2020.101513.
- Hill L, Prager Geller T, Baruah R, et al. Integration of a palliative approach into heart failure care: a European Society of Cardiology Heart Failure Association position paper. *Eur J Heart Fail* 2020; 22(12): 2327–2339, doi: 10.1002/ehf.1994.
- Wleklik M, Uchmanowicz I, Jankowska EA, et al. Multidimensional Approach to Frailty. *Front Psychol* 2020; 11: 564, doi: 10.3389/fpsyg.2020.00564.

Address for correspondence:

Maria Jędrzejczyk, MSc
Division of Research Methodology
Department of Nursing
Faculty of Nursing and Midwifery
Wrocław Medical University
5 K. Bartla Str
51-618 Wrocław
Tel.: 71 784 18 24
E-mail: maria.jedrzejczyk97@gmail.com

Received: 16.09.2024

Revised: 18.09.2024

Accepted: 28.10.2024